

# ALGEBRAIC RESULTS FOR CERTAIN VALUES OF THE JACOBI THETA-CONSTANT $\theta_3(\tau)$

Carsten Elsner and Yohei Tachiya

(submitted paper)

In its most elaborate form, the Jacobi theta function is defined for two complex variables  $z$  and  $\tau$  by  $\theta(z|\tau) = \sum_{\nu=-\infty}^{\infty} e^{\pi i \nu^2 \tau + 2\pi i \nu z}$ , which converges for all complex number  $z$ , and  $\tau$  in the upper half-plane. The special case

$$\theta_3(\tau) := \theta(0|\tau) = 1 + 2 \sum_{\nu=1}^{\infty} e^{\pi i \nu^2 \tau}$$

is called a Jacobi theta-constant or Thetanullwert of the Jacobi theta function  $\theta(z|\tau)$ . In this paper, we prove the algebraic independence results for the values of the Jacobi theta-constant  $\theta_3(\tau)$ . For example, the three values  $\theta_3(\tau)$ ,  $\theta_3(n\tau)$ , and  $D\theta_3(\tau)$  are algebraically independent over  $\mathbb{Q}$  for any  $\tau$  such that  $q = e^{i\pi\tau}$  is an algebraic number, where  $n \geq 2$  is an integer and  $D := (\pi i)^{-1} d/d\tau$  is a differential operator. This generalizes a result of the first author, who proved the algebraic independence of the two values  $\theta_3(\tau)$  and  $\theta_3(2^m\tau)$  for  $m \geq 1$ . As an application of our main theorem, the algebraic dependence over  $\mathbb{Q}$  of the three values  $\theta_3(\ell\tau)$ ,  $\theta_3(m\tau)$ , and  $\theta_3(n\tau)$  for integers  $\ell, m, n \geq 1$  is also presented.

**2010 MS Classification numbers:** 11J85, 11J91, 11F27

**Key Words:** Algebraic independence, Theta-constants, Nesterenko's theorem,  
Modular equations